

SUBGROUPS - I

OUT LINES

INTRODUCTION

DEFINITION- EXAMPLES

THEOREMS

ADDITIONAL INFORMATION

SUMMARY

PREPARED BY

S.Lt / Dr.D.CH.PAPARAO. M.Sc, Ph.D, PGDCA

Head, Dept. of Mathematics

S.K.B.R.COLLEGE, AMALAPURAM

East Godavari Dist. Andhra Pradesh

Mail ID: dchp876@gmail.com

INTRODUCTION

In Group Theory, a branch of Abstract Algebra, a Subgroup is a structure-preserving function in Groups.

COMPLEX:

Let G be a group and $H \neq \phi \subseteq G$ then H is called the Complex of G .

Any non-empty subset of a group G is called a complex.

Example:

1. $H = \{1\}$ is a complex of a multiplicative group $G = \{1, -1\}$
2. The set of even integers is a complex of $(\mathbb{Z}, +)$.

MULTIPLICATION OF COMPLEXES

Let H and K are two complexes of a group G then the product HK is defined as

$$\mathbf{HK} = \{ \mathbf{hk} / \mathbf{h} \in \mathbf{H}, \mathbf{k} \in \mathbf{K} \}$$

Note: 1. HK is also a complex of G .

2. Multiplication of Complexes is Associative.

$$\text{i.e., } H(KL) = (HK)L$$

INVERSE OF A COMPLEX

Let H be a complex of G , then the inverse of H

is defined as $H^{-1} = \{ h^{-1} / h \in H \}$

$$\text{i.e., } h \in H \Rightarrow h^{-1} \in H^{-1}$$

NOTE: We know that $(HK)^{-1} = K^{-1}H^{-1}$ and $(H^{-1})^{-1} = H$

SUBGROUP

Let (G, \cdot) be a group and $H \subseteq G$. H is said to be a Subgroup of G if H is itself a group w.r.t the binary operation ' \cdot ' of G .


EXAMPLES

1. $(\mathbf{Z}, +)$ and $(\mathbf{C}, +)$ are two groups w.r.t addition and $\mathbf{Z} \subseteq \mathbf{C} \Rightarrow (\mathbf{Z}, +)$ is a subgroup of $(\mathbf{C}, +)$.
2. $(\mathbf{Q}, +)$ and $(\mathbf{R}, +)$ are two groups w.r.t addition and $\mathbf{Q} \subseteq \mathbf{R} \Rightarrow (\mathbf{Q}, +)$ is a subgroup of $(\mathbf{R}, +)$.
3. The set of Even integers is a subgroup of $(\mathbf{Z}, +)$.

Relation between Group and Subgroup

Group: G

Subgroup: Subset & group



**Let G be group and H is a subgroup of G
 \Rightarrow 1. H is a subset of G . 2. H is a group.**

IMPROPER (OR) TRIVIAL SUBGROUPS

Let G be a group and $e \in G$. Then the sets $\{e\}$ and G are Subgroups of G . These two Subgroups are called Improper Subgroups of G .

All other subgroups other than $\{e\}$ and G are called PROPER or Non-Trivial Subgroups of G .

EXAMPLE

We know that $G = \{1, -1, i, -i\}$ is a multiplicative Group.

$H_1 = \{1\}$, $H_2 = \{1, -1\}$ and $H_3 = G = \{1, -1, i, -i\}$ are three Subgroups of G . (\because Identity element $e = 1$)

Clearly H_1 and H_3 are improper subgroups of G and H_2 is a proper subgroup of G .

Example for a Subgroup

Let $m \in \mathbf{Z}$ and $m\mathbf{Z} = \{ma / a \in \mathbf{Z}\}$ then the set $m\mathbf{Z}$ is a Subgroup of a group $(\mathbf{Z}, +)$.

Also $m\mathbf{Z}$ is cyclic Subgroup of $(\mathbf{Z}, +)$ generated by m .

$$\therefore m\mathbf{Z} = \{ma / a \in \mathbf{Z}\} = \langle m \rangle$$

Example for a Subgroup

Let X be a non empty subset and $S(X)$ be the set of all bijective mappings of X onto itself under the composition of mappings.

$$\therefore S(X) = \{f / f \text{ is a bijection on } X\}$$

For any $x_0 \in X$, we define $Hx_0 = \{f \in S(X) / f(x_0) = x_0\}$

then Hx_0 is a subgroup of $(S(X), \circ)$.

Q. What is the relation between between Complex and Subgroup ?

A. Eevery Subgroup is a Complex, but every Complex need not be a Subgroup.

Theorem : If H is a subgroup of a group G then $HH = H$

Proof : Let $x \in HH \Rightarrow x = h_1 h_2$, where $h_1 \in H, h_2 \in H$

$\therefore h_1 \in H, h_2 \in H \Rightarrow h_1 h_2 \in H \Rightarrow x \in H$

$\therefore x \in HH \Rightarrow x \in H \Rightarrow HH \subseteq H \rightarrow (1)$

Again $h \in H, e \in H \Rightarrow he \in HH \Rightarrow h \in HH$

$\therefore h \in H \Rightarrow h \in HH \Rightarrow H \subseteq HH \rightarrow (2)$

From (1) & (2), we get $HH \subseteq H$ & $H \subseteq HH \Rightarrow HH = H$

Hence proved

If H is a subgroup of a group G then $H^{-1} = H$

Proof : Let G be a group and $H \subseteq G$.

Part – I : Suppose H is a subgroup of G

Let $x \in H^{-1} \Rightarrow x = h^{-1}$, where $h \in H$

$\therefore h \in H \Rightarrow h^{-1} \in H$ (Inverse) $\Rightarrow x \in H$

$\therefore x \in H^{-1} \Rightarrow x \in H \Rightarrow H^{-1} \subseteq H \rightarrow (1)$

Part – II : Suppose $H^{-1} \subseteq H$

Let $h \in H \Rightarrow h^{-1} \in H \Rightarrow (h^{-1})^{-1} \in H^{-1}$

$\Rightarrow h \in H^{-1}$

$\therefore h \in H \Rightarrow h \in H^{-1} \Rightarrow H^{-1} \subseteq H \rightarrow (2)$

From (1) & (2) we get $H^{-1} = H$

Hence proved

**Note : The converse of the above theorem need not be true.
i.e., If $H = H^{-1}$ then H need not be a Subgroup of a group G .**

Consider a set $H = \{1\}$ and a multiplicate group $G = \{1, -1\}$

Clearly $H = H^{-1} = \{1\}$ and H is a Subgroup of G . $\because (e = 1)$

Again Consider a set $H = \{-1\}$ and a multiplicate group $G = \{1, -1\}$

Clearly $H = H^{-1} = \{-1\}$, but H is a not a Subgroup of G as

H has not a multiplicative identity $e = 1$. $\because H^{-1} = \{(-1)^{-1} = -1\}$

Let H be a Subgroup of a group G . The identity element of H is same as the identity element of G .

A nonempty subset H of a group G is a subgroup of G iff $a, b \in H \Rightarrow ab^{-1} \in H$

Proof: Suppose H be a Subgroup of a group G and e is the identity element of G .

Let $a, b \in H \Rightarrow a^{-1}, b^{-1} \in H \quad \because$ inverse axiom

For $a \in H, b^{-1} \in H \Rightarrow ab^{-1} \in H \quad \because$ identity axiom

$\therefore a, b \in H \Rightarrow ab^{-1} \in H$

Conversely suppose that $a, b \in H \Rightarrow ab^{-1} \in H \rightarrow (1)$

1. Identity axiom: Let $a \in H \Rightarrow a \in G \left[\because aa^{-1} = a^{-1}a = e \right]$

For $a, a \in H \Rightarrow aa^{-1} \in H \Rightarrow e \in H \quad \left[\text{using } (1) \right]$

where e is the identity element in H .

2. Inverse axiom: Let $a \in H, e \in h$ [$\because ae = ea = a$]

For $e, a \in H \Rightarrow ea^{-1} \in H \Rightarrow a^{-1} \in H$ [using (1)]

$\therefore a \in H \Rightarrow a^{-1} \in H$

3. Associative axiom: Let $a, b, c \in H \Rightarrow a, b, c \in G$

$\Rightarrow a(bc) = (ab)c \quad \because H \subseteq G$ & G is a group.

$\therefore '.'$ is associative in H .

4. Closure axiom: Let $a, b \in H \Rightarrow a^{-1}, b^{-1} \in H$

$\therefore a \in H, b^{-1} \in H \Rightarrow a(b^{-1})^{-1} \in H \Rightarrow ab \in H$

$\therefore '.'$ is a binary operation in H

$\therefore H$ is group and $H \subseteq G \Rightarrow H$ is a subgroup of G .

Hence proved

Necessary and Sufficient Condition for a Subgroup

Statement: A nonempty complex H of a group G is a subgroup of G if and only if

(i) $a, b \in H \Rightarrow ab \in H$, (ii) $a \in H \Rightarrow a^{-1} \in H$

Proof: Suppose H be a Subgroup of a group G
 $\Rightarrow H$ is itself a group w.r.t the operation of G

Let $a, b \in H \Rightarrow ab \in H \quad \because$ Closure axiom

For $a \in H \Rightarrow a^{-1} \in H \quad \because$ inverse axiom

Conversely suppose that

$a, b \in H \Rightarrow ab \in H \rightarrow (1)$ and $a \in H \Rightarrow a^{-1} \in H \rightarrow (2)$

1. Closure axiom: Let $a, b \in H \Rightarrow ab \in H$ [using (1)]

2. Associative axiom: Let $a, b, c \in H \Rightarrow a, b, c \in G$
 $\Rightarrow a(bc) = (ab)c \quad \because G$ is a group & $H \subseteq G$

3. Existence of inverse: Let $a \in H \Rightarrow a^{-1} \in H$ (\because using 2)

\therefore Every element in H having multiplicative inverse in H .

4. Identity axiom: Let $a, b \in H \Rightarrow a^{-1}, b^{-1} \in H$ (\because using 2)

For $a, a^{-1} \in H \Rightarrow aa^{-1} \in H \Rightarrow e \in H$ ($\because aa^{-1} = a^{-1}a = e$)

\Rightarrow 'e' is a the identity element in H .

$\therefore (H, \cdot)$ is a group and $H \subseteq G \Rightarrow H$ is a Subgroup of G .

Hence proved

NOTE :

1. Let H be a complex of an multiplicative group (G, \cdot) .

Then H is a Subgroup of (G, \cdot) iff $a, b \in H \Rightarrow ab^{-1} \in H$.

2. Let H be a complex of an additive group $(G, +)$.

Then H is a Subgroup of $(G, +)$ iff $a, b \in H \Rightarrow a - b \in H$.

S.T $m\mathbb{Z}$ is a subgroup of \mathbb{Z} , where m is fixed + ve integer.

Sol: Let \mathbb{Z} be the set of all integers and m is fixed + ve integer.

Let $x, y \in m\mathbb{Z}$, where $m\mathbb{Z} = \{ma / a \in \mathbb{Z}\}$

$\Rightarrow x = ma$ and $y = mb$ for $a, b \in \mathbb{Z}$

Now $x - y = ma - mb = m(a - b) = ma^1 \in m\mathbb{Z} \quad \because a^1 = (a - b) \in \mathbb{Z}$

$\Rightarrow m\mathbb{Z}$ is a subgroup of $(\mathbb{Z}, +)$.

Note : In general $3\mathbb{Z}, 4\mathbb{Z}, 5\mathbb{Z}, \dots$ etc are subgroups of $(\mathbb{Z}, +)$.

Example Problem:

Prove that the set of multiples of 3 is a subgroup of the group of integers under addition.

Here, the set of multiples of 3 = $3\mathbb{Z} = \{3a / a \in \mathbb{Z}\}$

Union of two subgroups need not be a Subgroup.

We know that $2\mathbb{Z}$ and $3\mathbb{Z}$ are two Subgroups of $(\mathbb{Z}, +)$.

$$\because 2\mathbb{Z} = \{\dots -4, -2, 0, 2, 4, \dots\} \ \& \ 3\mathbb{Z} = \{\dots -6, -3, 0, 3, 6, \dots\}$$

$$\Rightarrow \text{Let } H = 2\mathbb{Z} \cup 3\mathbb{Z} = \{\dots -6, -4, -3, -2, 0, 2, 3, 4, 6, \dots\}$$

Let $2, 3 \in H \Rightarrow 2 - 3 = -1 \notin H \Rightarrow H$ is not a subgroup of $(\mathbb{Z}, +)$.

Intersection of two subgroups is also a Subgroup.

Give an example of a nonabelian group

Example: The $G = \{r_0, r_1, r_2, f_1, f_2, f_3\}$ of all symmetries of an equilateral triangle forms a nonabelian group w.r.t to the composition of mappings 'o' on G.

1. Can an abelian group have a nonabelian Subgroup?

Ans: NO \Rightarrow Every subgroup of an abelian is abelian.

2. Can a nonabelian group have an abelian Subgroup?

Ans: YES

Example-1: The set $\{e\}$ is a subgroup of a nonabelian group G , where e is the identity element of G .

Example-2: Let G be a nonabelian group and $x \in G$, $x \neq e$.

Then $\langle x \rangle$ is an abelian subgroup of G . For example, S_3 is a nonabelian group such that only the cyclic subgroups are abelian. where S_3 is the permutation group on 3 symbols.

3. Can a nonabelian group have a nonabelian Subgroup?

Ans: YES

Note: The dihedral groups are examples of non – abelian groups

Every subgroup of an abelian group is abelian

Let G be an abelian group and H is a subgroup of G .
We prove that H is an abelian group

Let $a, b \in H \Rightarrow a, b \in G \quad (\because H \subseteq G)$

$\Rightarrow ab = ba \quad (\because G \text{ is abelian group})$

$\therefore H$ is a subgroup of G .

Find all subgroups of the group $(G = \{0, 1, 2, 3, 4, 5\}, \oplus_6)$

$H_1 = \{0\}, H_2 = \{0, 3\}, H_3 = \{0, 2, 4\}, H_4 = \{0, 1, 2, 3, 4, 5\} = G$

Here H_1 and H_4 are improper subgroups and H_2 and H_3 are proper subgroups of G .

Intersection of two subgroups of a group is also a subgroup

Proof: Let H and K are two subgroups of a group G .

Claim: We prove that $H \cap K$ is a subgroup of G

Let $a, b \in H \cap K \Rightarrow a, b \in H$ and $a, b \in K$

For $a, b \in H \Rightarrow ab^{-1} \in H \quad \because H$ is a subgroup of G

For $a, b \in K \Rightarrow ab^{-1} \in K \quad \because K$ is a subgroup of G

Using above, $ab^{-1} \in H$ and $ab^{-1} \in K \Rightarrow ab^{-1} \in H \cap K$

$\therefore H \cap K$ is a subgroup of G

NOTE:1. H is a subgroup of a group G iff $a, b \in H \Rightarrow ab^{-1} \in H$

2. The intersection of any family of a subgroups of a group G is also a subgroup of G .

Let H and K be two subgroups of a group G . Then prove that $H \cup K$ is a subgroup of G iff $H \subseteq K$ or $K \subseteq H$

Proof: Let H and K be two subgroups of a group G .

PART-I: Suppose that $H \cup K$ is a subgroup of G

We prove that $H \subseteq K$ or $K \subseteq H$

If possible suppose that $H \not\subseteq K$ and $K \not\subseteq H$

$\therefore H \not\subseteq K$, then there exists $a \in H$ and $a \notin K \rightarrow (1)$

$\therefore K \not\subseteq H$, then there exists $b \in K$ and $b \notin H \rightarrow (2)$

From (1) & (2), $a \in H \Rightarrow a \in H \cup K$, $b \in K \Rightarrow b \in H \cup K$

$\therefore a \in H \cup K$, $b \in H \cup K \Rightarrow ab \in H \cup K$ (Closure)

$\Rightarrow ab \in H$ or $ab \in K$ or $ab \in H \cap K$

Case -I : Suppose that $ab \in H$

$\therefore a \in H \Rightarrow a^{-1} \in H$ and $ab \in H \Rightarrow a^{-1}(ab) \in H \Rightarrow b \in H$

It is contradiction (\because from (2), $b \notin H$)

\therefore Our supposition $ab \in H$ is wrong. i.e., $ab \notin H \rightarrow (3)$

Case -II : Suppose that $ab \in K$

$\therefore b \in K \Rightarrow b^{-1} \in K$ and $ab \in K \Rightarrow (ab)b^{-1} \in K \Rightarrow a \in K$

It is contradiction (\because from (1), $a \notin K$)

\therefore Our supposition $ab \in K$ is wrong. i.e., $ab \notin K \rightarrow (4)$

From (3) & (4), $ab \notin H, ab \notin K \Rightarrow ab \notin H \cup K$ (Closure failed)

Hence our supposition $H \not\subseteq K$ and $K \not\subseteq H$ is false.

$\therefore H \subseteq K$ or $K \subseteq H$.

PART - II : Suppose that $H \subseteq K$ or $K \subseteq H$.

$\therefore H \subseteq K \Rightarrow K = H \cup K \Rightarrow H \cup K$ is a subgroup of G .

$\therefore K \subseteq H \Rightarrow H = H \cup K \Rightarrow H \cup K$ is a subgroup of G .

Hence proved

Let H and K be two subgroups of a group G.

Then prove that HK is a subgroup of G iff $HK = KH$

Proof: Let H and K be two subgroups of a group G.

\because H is a subgroup of G $\Rightarrow H = H^{-1}$ and $HH^{-1} = H \rightarrow$ (1)

\because K is a subgroup of G $\Rightarrow K = K^{-1}$ and $KK^{-1} = K \rightarrow$ (2)

PART – I: Suppose HK is a subgroup of G.

\because HK is a subgroup of G $\Rightarrow (HK)^{-1} = HK$

$$\Rightarrow (K^{-1}H^{-1}) = HK$$

$$\Rightarrow KH = HK$$

PART-II: Suppose $HK=KH$

Now $(HK)(HK)^{-1} = (HK)(K^{-1}H^{-1}) = H(KK^{-1})H^{-1} = (HK)H^{-1}$

$$= (KH)H^{-1} = K(HH^{-1}) = KH = HK$$

\therefore HK is a subgroup of G.

If H is a subgroup of a group G then $HH^{-1} \subseteq H$

Proof : Let G be a group and $H \subseteq G$.

Suppose H is a subgroup of G

Claim : We prove that $HH^{-1} \subseteq H$

Let $x \in HH^{-1} \Rightarrow x = ab^{-1}$, where $a \in H, b^{-1} \in H^{-1}$

$\because b^{-1} \in H^{-1} \Rightarrow b \in H \Rightarrow b^{-1} \in H$ ($\because H$ is a subgroup)

$\therefore a \in H, b^{-1} \in H \Rightarrow ab^{-1} \in H \Rightarrow x \in H$

$\therefore x \in HH^{-1} \Rightarrow x \in H \Rightarrow HH^{-1} \subseteq H$

Hence proved

ADDITIONAL INFORMATION

The following web links are very useful to develop the subject skills and learning the problem solving techniques in easy manner.

<https://web.ma.utexas.edu/users/rodin/343K/Subgroups.pdf>

<http://www.maths.lth.se/matematiklth/personal/ssilvest/AlgebraVT2011/Lecture09Silvestrov.pdf>

<https://www.youtube.com/watch?v=TJAQNIGvfjE>

<https://www.youtube.com/watch?v=dpj2dV0scl0>

<http://web.math.princeton.edu/swim/SWIM%202010/SWIM%202010%20Course%20II%20Lecture%20Notes%206%20of%207.pdf>

<https://www.youtube.com/watch?v=aWxzn00MPsk>

SUMMARY

I have concluded that PPT presentation is very useful in establishing objectives, Basic Concepts and illustrating concrete examples.

I hope that utilizing all of these methods through PPT Slides helps to engage students with different types of learning styles.

I have added definitions, example problems and theorems of the chapter Subgroups in brief and short methods.

S.K.B.R.COLLEGE AMALAPURAM

